Quantification of DNA double strand breaks and oxidation response in children and adults undergoing dental CBCT scan.

1	Authors					
2	Niels Belmans ^{1,2} , Liese Gilles ¹ , Randy Vermeesen ² , Piroska Virag ³ , Mihaela Hedesiu ⁴ , Benjamin Salmon ⁵ ,					
3	Sarah Baatout ² , Stéphane Lucas ⁶ , Ivo Lambrichts ¹ , Reinhilde Jacobs ^{7,8} , Marjan Moreels ^{2*} & DIMITRA					
4	Research Group					
5						
6	¹ Morphology Group, Biomedical Research Institute, Hasselt University, Agoralaan Building C, Diepenbeek, Belgium					
7	² Belgian Nuclear Research Centre, Radiobiology Unit, SCK•CEN, Mol, Belgium					
8	³ Institute of Oncology "Prof. dr. Ion Chiricuta", Cluj-Napoca, Romania					
9	⁴ 'Iuliu Hatieganu' University of Medicine and Pharmacy, Department of Oral and Maxillofacial Radiology, , Cluj-Napoca, Romania					
10	⁵ Paris Descartes University - Sorbonne Paris Cité, EA 2496 - Orofacial Pathologies, Imaging and Biotherapies Lab and Dental					
11	Medicine Department, Bretonneau Hospital, HUPNVS, AP-HP, Paris, France					
12	⁶ Namur Research Institute for Life Sciences, University of Namur, Namur, Belgium					
13	⁷ Katholieke Universiteit Leuven, Department of Imaging and Pathology, OMFS-IMPATH Research group, and University					
14	Hospitals, Oral and Maxillofacial Surgery, Dentomaxillofacial Imaging Center, Kapucijnenvoer 7, Leuven, Belgium					
15	⁸ Karolinska Institutet, Department Dental Medicine, Huddinge, Sweden					
16						
17	*: Corresponding author:					
18	Dr. Marjan Moreels					
19	Belgian Nuclear Research Centre					
20	Institute for Environment, Health and Safety – Interdisciplinary Biosciences – Radiobiology Unit					
21	Boeretang 200					
22	2400 Mol					
23	marjan.moreels@sckcen.be					

24

+32 14 33 28 16

25 Supplementary Table 1

26 27

28

29

30 31


32

Supplementary Table 1. Power analysis for one-way repeated analysis of variance and pair t-tests.

	Statistical test	N	Effect size (F or Cohen)**	Significance level	Power
DNA damage: children	One-way RM ANOVA*	38	2.2	.05	1
DNA damage: adults	One-way RM ANOVA	13	2.2	.05	.9
8-oxo-dG: children	Paired t-test (2-sided)	68	.666	.05	.99
8-oxo-dG: adults	Paired t-test (2-sided)	19	0.833	.05	.93
FRAP: children	Paired t-test (2-sided)	117	1.888	.05	1
FRAP: adults	Paired t-test (2-sided)	17	1.888	.05	.99

^{*:} RM ANOVA = repeated measures analysis of variance

Supplementary Data 1




Supplementary Data 1. Absorbed radiation dose does not only depend on the device uses, but also on the field of view (FOV) and scanning protocol used during the examination. (left). Patients examined using a Promax 3D device receive on average a higher radiation dose than those examined with a Accuitomo 170 device or NewTom device. However, these data do not take into account the FOV or the scanning protocol. (right). Radiation dose increases with increasing FOV and resolution of the scan. This is seen for all devices (except for Accuitomo 170 for which only one scanning protocol was used). Furthermore, the radiation dose is higher when high resolution (HiRes) protocols were used. This explains the differences seen in the left panel, since for Planmeca ProMax and NewTom HiRes protocols were used, whereas only standard protocols were used in Accuitomo. No significances were shown in the right panel. ***: p < .0002; ****: p < .0001.

^{**:} F value for one-way RM ANOVA and Cohen's effect size for paired t-tests. 58,59

34 Supplementary Data 2

35 36



Supplementary Data 2. Comparison of % change in 8-oxo-dG excretion shows no difference between boys and girls. The proportional change in 8-oxo-dG levels does not differ between boys and girls (Mann-Whitney U value = 431, p = .203). Green dotted line = average.